



Editor: Alan N. Moshell, M.D.

Skin-Nervous System Interactions

John C. Ansel, Alan H. Kaynard,* Cheryl A. Armstrong, John Olerud,† Nigel Bunnett,‡ and Donald Payan§

Department of Dermatology, Emory University School of Medicine, Atlanta, Georgia; *Dermatology Service, VA Medical Center, Portland, Oregon; †Department of Dermatology, University of Washington, Seattle, Washington; ‡Department of Physiology and Surgery, University of California San Francisco, San Francisco; and §Kepri Pharmaceuticals, South San Francisco, California, U.S.A.

Components of the cutaneous nervous system interact with multiple types of cells in the skin to mediate actions important in skin inflammation and wound healing [1-5]. Over 100 years ago it was observed that antidromic stimulation of the sensory nerves emanating from dorsal root ganglia produced vasodilation and apparent local inflammation of the skin (for review see [6]). This observation, repeated many times since, led to the understanding that the cutaneous sensory nervous system (CSNS) not only relays sensory information from the skin to the central nervous system (CNS), but also plays an effector role in the skin's inflammatory response [7-9].

Sensory neurons express at least 17 different neuropeptides [6]. Several of these, substance P (SP) [6,10,11], calcitonin gene-related peptide (CGRP) [12-14], substance K (SK) [10,15,16], and vasoactive intestinal polypeptide [17-20], have activities or cellular localizations that suggest involvement in tissue repair. SP is the prototype of neuropeptides released from sensory C-fibers in the skin [21]. The activities of SP require not only secretion but also the expression of the SP receptor (SPR) on local target cells and expression of tissue proteases that degrade neuropeptides, such as neutral endopeptidase (NEP).

Another important facet of nervous system-cutaneous interactions are neurotrophic growth factors that mediate cutaneous reinnervation, e.g., nerve growth factor (NGF) [22,23], brain-derived nerve growth factor [24,25], neurotrophin-3 [26], and human neurotrophin-5 [27]. NGF is produced by the neural Schwann cells and by the regenerating epidermis itself [28,29].

Reprint requests to: Dr. John C. Ansel, Emory University School of Medicine, Department of Dermatology, 5311 Woodruff Memorial Building, Atlanta, GA 30322.

Abbreviations: CGRP, calcitonin gene-related peptide; CNS, central nervous system; CSNS, cutaneous sensory nervous system; NEP, neutral endopeptidase; SK, substance K; SP, substance P; SPR, substance P receptor.

BIOLOGIC ACTIVITIES OF SP AND SPR

SP Biochemistry and Tissue Distribution SP, an 11-amino acid peptide that is a member of the tachykinin family of neuropeptides, is present in many areas of the nervous system but is one of a few neuropeptides restricted to cells of neural crest origin [30,31]. It is located in nerve endings throughout the body, but especially in areas of immunologic importance such as the skin, gastrointestinal tract, respiratory tract, and eye [31]. In the peripheral nervous system, SP occurs in a subpopulation of primary afferent neurons including A- δ pain fibers and C-fibers, which transmit impulses initiated by noxious stimuli, and in nerve plexuses of the gastrointestinal tract. The role of SP as a neurotransmitter is supported by its presence in synaptosomal fractions and large granular vesicles in nerve endings of synapses within dorsal horn cells [32].

SP is synthesized in the dorsal root ganglia, from which it migrates centrally to the dorsal horn of the spinal cord and peripherally to nerve terminals of sensory neurons [21]. A single gene has been identified that can be differentially processed into two different SP precursor mRNAs, α - and β -preprotachykinin [33,34]. The first contains the sequence of SP only whereas the second contains the sequences of SP as well as a second tachykinin-peptide, SK [33,35]. SP and SK share a common spectrum of biologic activities [10,36,37], but levels of the two precursor mRNAs vary between tissues [33,38], the CNS producing more α -preprotachykinin and the peripheral nervous system (PNS) producing more β -preprotachykinin [34]. SP is degraded by a specific thiorphan-sensitive enkephalinase, CD10, also known as NEP [39-41].

SP Receptor Biochemistry The amino acid sequence of the human and rat SPR neurokinin (NK₁), has been deduced from molecular cloning data [42-44]. The receptor protein is composed of 407 residues with seven putative membrane-spanning domains and shares sequence homology with the SK receptor (NK₂) in its transmembrane regions and in its first and second cytoplasmic loops [45,46]. The SPR mRNA is expressed by neurons and glia in the

Table I. *In Vitro* Effects of the Neuropeptide Substance P on the Immune System

Effect	Specifics
Cellular proliferation	● Enhances T-cell proliferation (much greater in Peyer's patches than in spleen) [98]
Ig production	● Increase IgA and IgM [102] ● Increase heavy-chain mRNA [102]
Cytokine production/secretion	● Enhances IL-2 synthesis and secretion by peripheral blood monocyte cells and HUT 78 cells [105] ● Enhances IL-2 receptor expression on Jurkat-SP-R cells [104] ● Enhances IL-1, IL-6, TNF α , and interferon γ production by monocytes [103] ● Enhances expression and secretion of IL-1 α , IL-1 β , and granulocyte/macrophage colony-stimulating factor by keratinocytes [57] ● Enhances TNF α secretion by mast cells [56]
Phagocytosis	● Enhances macrophage and polymorphonuclear-leukocyte activity [108,185,186]
Mediator release	● Enhances histamine, prostaglandin, and leukotriene release from mast cells [106]
Cell homing, chemotaxis	● Enhances human monocyte chemotaxis [186,187] ● Alters CD2 and CD45 levels on Jurkat-SP-R cells [104,188]

CNS, neurons of the myenteric plexus, smooth muscle cells, acinar cells, endothelial cells, fibroblasts, various types of circulating immune cells, and skin [42,47–50]. Sites of expression agree well with earlier autoradiographic localizations [51–53]. SP, with a kD of 0.5–1.0 nM, is the most potent ligand for the NK₁ receptor [54]. Activation of the NK₁ receptor results in inositol phospholipid hydrolysis and the transient appearance of inositol 1,4,5 triphosphate, the release of Ca⁺⁺ from internal stores, and an oscillatory chloride current [55]. We have identified SPR in the skin on keratinocytes and mast cells [56,57], whereas other recent studies have demonstrated SPR on endothelial cells [58,59].

Actions of SP In the CNS, SP is thought to act in the cortex, hypothalamus, and lower brainstem [60–67]. SP in the PNS has numerous actions including contraction of the smooth muscle layers in the gastrointestinal tract [68,69]; production of edema and bronchoconstriction in the respiratory tract [70–72]; enhancement of nasal secretion [70,73]; stimulation of tracheal, epithelial, and glandular secretion of glycoprotein-rich fluid [74]; calcium-dependent secretion by the parotid and submandibular glands [75]; and modulation of both exocrine and endocrine pancreatic function [76–79]. Blood and tissue concentrations [80–82] of SP suggest that the peptide is available locally at functionally relevant concentrations. The peripheral actions of SP possibly involved in cutaneous neuroinflammation fall into three main categories: 1) vasodilation and increased vascular permeability, 2) local inflammation and effects on the immune system, and 3) promotion of cellular proliferation.

Vasodilation and Increased Vascular Permeability Increased blood flow and vascular permeability are effects well established in several animal species and humans [83–85]. Vasodilation is due to direct actions of SP on vascular smooth muscle [85,86] and enhanced production of nitric oxide by the endothelium [87]. SP is also the mediator of antidromic vasodilation, as shown by its presence in unmyelinated primary sensory fibers in the skin [86,88], its release from peripheral endings of sensory neurons during antidromic stimulation [89,90], and its vasodilatory effects after venous infusion [91]. In addition, SP can initiate microvascular permeability and protein extravasation after tissue injury [31,91,92]. Intradermal injection of SP in humans produces a wheal caused by increased vascular permeability followed by a flare reaction caused by reflex neurogenic vasodilation [93–95]. The reflex nature of the flare is demonstrated by its inhibition by local anesthetics; the wheal response remains unaffected [96,97].

Local Inflammation and Specific Effects on the Immune System SP has been shown to have stimulatory effects on the immune system both *in vitro* and *in vivo* [98–100] (Table I). Initially, SP was found in high concentrations in inflamed tissues such as the joints in experimental arthritis in the rat [101]. Later, injections of SP into normal joints produced inflammation that mimicked adjuvant-induced arthritis [101].

Nanomolar concentrations of SP enhance proliferation of both

human and murine peripheral-blood T lymphocytes [98,99]. [³H]-thymidine and [³H]-leucine uptake by human T lymphocytes is stimulated up to 60–70% by both SP and a C-terminal fragment of SP, SP_{4–11}, both in the presence and absence of other mitogens [98]. SP also increases, by up to 300%, the synthesis of immunoglobulins from mixed lymphocyte cultures from the spleen, lymph nodes, and Peyer's patches, the major effect being on IgA [99]. More recently, Pascual *et al.*, using an IgM- and IgA-producing B-lymphoma cell line, showed that SP can synergize with lipopolysaccharide to increase the production of these two immunoglobulins [102]. Continuous infusion of SP *via* miniosmotic pumps stimulated proliferation of lymphocytes isolated from the spleen and Peyer's patches in mice, and immunoglobulin synthesis, particularly IgA, was also increased [100]. SP has effects on cytokine regulation including enhanced production of cytokines such as interleukin 1 (IL-1), IL-6, tumor necrosis factor α (TNF α), and interferon- γ by monocytes [103], increase in IL-2 receptors in a Jurkat T-cell line expressing a transfected SPR cDNA [104], and augmented IL-2 synthesis in peripheral blood monocyte cells and HUT 78 cells [105].

SP can also modulate acute hypersensitivity responses. At micromolar concentrations SP evokes the release of histamine from rat serosal or connective-tissue-type mast cells [106]. Unlike other basic peptides, which stimulate mast cells nonspecifically, SP action is cell specific and includes stimulation of the generation of unstored mediators such as leukotrienes [107,108]. The mast cell responses elicited by SP were distinguished from those dependent on IgE by the apparent lack of a requirement for extracellular calcium [106].

We have recently demonstrated [56] that mast cell TNF α mRNA is selectively up-regulated by SP in a concentration-dependent manner. SP increased secreted TNF α from both cloned murine CFTL 12 mast cells and freshly isolated peritoneal mast cells. The addition of SP to cultured murine and normal human foreskin keratinocytes resulted in concentration-dependent increases in mRNA for IL-1 α , IL-1 β , and IL-1 α that were maximal after 3 h of exposure ([57], unpublished observations), but no effects on keratinocyte TNF α or IL-8 were found. It is interesting that SP_{1–9}, but not the C-terminal fragments SP_{4–11} or SP_{7–11}, are capable of inducing production of some cytokines by keratinocytes. More recently, we observed that SP can also specifically activate normal human microvascular endothelial cells to produce the neutrophil activating cytokine IL-8.¶

Promotion of Cell Proliferation Neuropeptides, especially SP, have also been implicated in the modulation of inflammation and wound healing by the ability to promote proliferation of various types of target cells. SP has been shown to act through the

¶ Kramp J, Brown J, Cook P, Russell B, Lawley T, Armstrong C, Ansel J: Neuropeptide induction of human microvascular endothelial cell interleukin 8 (abstr). *J Invest Dermatol* 104:586, 1995.

SPR to stimulate proliferation of cultured keratinocytes [11]. Additionally, SP and SK can stimulate proliferation of arterial smooth muscle cells and therefore may assist in angiogenesis during healing [10,16,109]; SP has also been shown to stimulate neovascularization *in vivo* [110]. SP and SK stimulate the proliferation of both human skin fibroblasts [10] and cultured endothelial cells [110], suggesting that SP released from the skin in association with tissue injury may not only enhance vasodilation and the inflammatory response, but may stimulate proliferation of cutaneous epithelial, vascular, and connective tissue.

ROLE OF NEP IN REGULATING NEUROPEPTIDE ACTIVITY

NEP Biochemistry and Tissue Distribution The biologic actions of neurotransmitters such as acetylcholine are terminated by enzymatic degradation, by re-uptake into nerve endings, and/or by diffusion away from the target cell [111,112]. Accumulating evidence suggests that neuropeptides are degraded and inactivated by cell surface peptidases [113]. The best-studied cell surface peptidase, neutral endopeptidase (EC 3.4.24.11, NEP), also known as enkephalinase, common acute lymphoblastic leukemia antigen, or CD 10, is the major tachykinin-degrading enzyme. NEP was discovered in the brush border of the kidney in 1968 as an enzyme that hydrolyzed the β chain of insulin [114] but was not obtained in pure form until 1974 [115,116] and has also been isolated from other tissues. Interest in NEP was rekindled in 1978 when NEP was shown to degrade enkephalins in the brain [117]. NEP degrades tachykinins, opioid peptides, GRP-10, somatostatin-14, neurotensin, cholecystokinin (CCK), gastrin, vasoactive intestinal polypeptide, CGRP, and others [118–120]. Although tachykinins and enkephalins are the best substrates, the proximity of peptidergic neurons to cells that express NEP also determines whether a neuropeptide can be a relevant physiologic substrate for NEP. The consensus is that NEP in the nervous system is found in neurons rather than glial cells. In the dorsal horn of the spinal cord and the trigeminal nucleus, staining for NEP, SP, and enkephalins is superimposable.

The search for a physiologic role of NEP has been facilitated by use of the specific inhibitors phosphoramidon and thiorphan. The release of both enkephalins and SP from brain slices *in vitro* is enhanced in the presence of NEP inhibitors [121,122], and these inhibitors relieve pain [122–124], which is consistent with their effects on endogenous opioid peptide release and stability. In the airways, inhibitors of NEP potentiate effects of SP on tracheal secretion [74], bronchomotor tone [125], and cough [126]. In addition, inhibition of NEP has been shown to potentiate tracheal neurogenic inflammation [127]. The mechanism by which NEP modulates the effects of neuropeptides is widespread and phylogenetically ancient. For example, phosphoramidon stimulates the growth of small-cell lung carcinoma cells by preventing the inactivation of endogenous GRP (an autocrine growth factor) by NEP [128]. An NEP-like enzyme is also found in hemocytes of the mollusc *Mytilus edulis*, and inhibition of NEP activity decreases by five orders of magnitude the amount of Met-enkephalin required to activate these cells [129].

Although regulation of NEP activity and expression has received little attention, changes in its activity have profound effects on the cellular response to neuropeptides. NEP activity in the trachea is reduced by up to 50% by infection [130], and NEP is down-regulated by intestinal inflammation; as a result, the degradation of SP by inflamed tissue is markedly reduced [131]. The ability of glucocorticoids to stimulate expression of NEP by transformed tracheal epithelial cells [132] and to down-regulate SPR [133] may partly explain their anti-inflammatory properties.

CUTANEOUS NGF

NGF Biochemistry and Tissue Distribution NGF, a 118-amino acid, ~130-kD protein isolated more than 40 years ago as an agent stimulating growth and neurite development in embryonic sensory cells [134], is the best-characterized member of the neuro-

trophin family [135]. NGF is composed of three subunits, α , β , and γ , but biologic activity appears to reside in the 26-kD β subunit (2.5S NGF) [136].

In the mature animal, NGF is expressed in the CNS and peripheral tissues [137,138] including keratinocytes of mouse and human epidermis [28,29,139]. NGF mRNA in keratinocytes is increased by ultraviolet radiation and phorbol esters, and reduced by hydrocortisone [28,29]. Anti-NGF monoclonal antibody stained only basal keratinocytes obtained from growing colonies; cells from confluent cultures and fully differentiated cells showed no evidence of NGF staining [29], suggesting a role for NGF production by the regenerative epidermis in wound healing.

NGF Receptor One NGF receptor (NGFR) protein is a more abundant low-molecular-weight protein of ~75 kD, p75^{NGF-R}, that binds NGF with low affinity, and the second is a ~140-kD protein that binds NGF with high affinity [140]. Both are required for high-affinity binding and a normal cellular response. The transmembrane protein p75^{NGF-R} appears to serve as a common subunit for different neurotrophin receptors, whereas the specificity of the receptor is conferred by the second subunit [141]. The higher-molecular-weight subunit is now known to be the proto-oncogene, *trk*, a member of the tyrosine kinase receptor family [142,143].

General Biologic Actions of NGF NGF has been shown to be both neurotrophic and neurotactic in the CNS. The outgrowth of processes from ganglion neurons is dependent on NGF [144], and NGF is also crucial for neuronal survival and function. Exposure of fetal rats to antibodies to NGF causes the ablation of an entire population of sensory neurons including those containing SP [145,146]. NGF is taken up by sensory neurons and transported in retrograde fashion to the cell body [147]. Deprivation of NGF results in reduced levels of SP in dorsal root ganglia [148].

Effects of Wounding on Skin Innervation Innervation of skin is altered by wounding [149–153]. In rat and guinea pig skin, SP-containing nerve fibers disappear from wounded skin within 1–2 d [152,153] and regenerating fibers begin to return after 3 d, increasing gradually for ~2 weeks after wounding and resulting in hyperinnervation. The new fibers exhibit extensive sprouting and form an unusually dense network just beneath the regrowing epidermis and then decrease in number over several weeks, following the course of healing and suggesting a functional link between regeneration of the CSNS and the integrity of the skin.

Role of NGF in Reinnervation of Wounded Skin After injury to a cutaneous nerve, denervated skin is reinnervated by two mechanisms, axonal regeneration (i.e., the regrowth of injured axons) and collateral reinnervation (i.e., the sprouting of undamaged axons in adjacent tissues into the denervated area) [154]. The latter has been shown to be completely dependent on NGF, because reinnervation does not occur after depletion of NGF [155]. Furthermore, collateral reinnervation is blocked by passive or active immunization against NGF [23,156]. Diamond *et al* went on to show that intradermal injection of NGF produced collateral sprouting in normal, fully innervated skin, increased the rate of sprouting in denervated skin, and restored the rate of sprouting in anti-NGF-treated denervated skin [23]. Finally, there appears to be an increase in synthesis of NGF within denervated tissue [23,157]. As mentioned above, keratinocytes express and synthesize NGF and may be the source of the endogenous growth factor utilized in sprouting. This is especially likely because a) reinnervating nerves enter the wound from the edge along with the epithelial tongue [152] and b) there is an unusually dense arborization of nerves just beneath the rapidly growing keratinocyte layer, which contains keratinocytes known to express the highest level of NGF [29]. It is possible, however, that NGF from Schwann cells, fibroblasts, and smooth muscle cells may also be involved.

NGF may also play a significant role in axonal regeneration after injury [158]. This hypothesis is based upon several observations: a) dorsal root ganglia have high-affinity NGF receptors [140]; b)

Schwann cells make NGF [159]; c) NGF and NGFR levels increase in degenerating nerve pathways and then become down-regulated when regenerating axons arrive [160]; and d) when peripheral nerves are cut and begin to degenerate, their Schwann cells start to express low-affinity NGFR and this expression disappears when the regenerating axon arrives [158,160]. It is thought that NGF is bound to the NGFR on Schwann cells and is then transferred to high-affinity NGFR on the regenerating axons because of the differences in binding affinities of the two types of receptor. The role of NGF in axonal regeneration has been questioned recently by evidence that nerve regeneration is unaffected by treatment with anti-NGF antibodies [161].

Proinflammatory and Proliferative Actions of NGF Paracrine actions of NGF may also contribute to the maintenance of the integrity of skin. NGF causes proliferation and release of histamine from mast cells [162], promotes differentiation of specific granulocytes, enhances T-lymphocyte-dependent antibody synthesis, and induces growth and differentiation of human B lymphocytes [163,164]. Furthermore, NGF may modulate melanocyte gene expression, is chemotactic for melanocytes, and can enhance melanocyte dendricity [139].

ROLE OF THE NEUROLOGIC SYSTEM IN WOUND HEALING

Effects of Denervation on Wound Repair It is also an old observation that chronic wounds and ulcers occur in skin with absent or dysfunctional innervation, an event often attributed to pressure, ischemia, infection, or incidental trauma related to a lack of sensation. Once such wounds were formed, failure to heal was attributed to difficulty with hygiene, protection of the wound from additional trauma, worsening of the underlying disease, poor nutrition, and susceptibility to infection. As early as 1927 it was noted that damage to the PNS is capable of altering growth and repair of skin, leading ultimately to ulceration and necrosis of the affected area [165]. Such observations led to the hypothesis that cutaneous innervation provides "trophic" support to skin integrity. Interest in this phenomenon has focused primarily on the role of sensory nerves, because it was noted that the flare response, a reflex of the CSNS [166,167], is absent in denervated skin that develops chronic wounds. Additional information has subsequently implicated the CSNS as a major contributor to skin integrity. For example, diseases and injuries that produce CSNS deficits (e.g., diabetes [168], herpes zoster [169], and spinal cord injury [170,171]) are often characterized by poor wound healing.

Denervation of rabbit skin by cutting of the nerves impairs the healing of incisional wounds [172]. Scars in denervated skin showed decreased tensile strength, hydroxyproline content, and protein composition compared to those in control skin. Near total sensory denervation can also be accomplished by treatment with high doses of capsaicin [173]. Kjartansson *et al* studied the effects of capsaicin-induced sensory denervation on the survival of tissue in experimental critical skin flaps [174]. Ten days after surgery, only 32% as much tissue remained in flaps of denervated skin as compared to flaps of innervated skin. Severe corneal lesions with opacity, edema, and ulceration can be produced in neonatal mice and rats by ablation of sensory neurons with capsaicin [175]. Similar results can be obtained by cutting the trigeminal nerve [176]. In a related experiment, depletion of neuropeptides in rabbit corneas by topical or retrobulbar injection of capsaicin caused a delay in corneal wound healing [177]. Such treatments reduce the migration rate of epithelial cells surrounding an n-heptanol-induced epithelial wound. These findings support the hypothesis that the trophic effects of sensory nerves on corneal or epidermal epithelium are mediated by neuropeptides in the sensory nerve endings.

Changes in Tissue Neuropeptide Levels After Wounding The few studies that have been conducted regarding neuropeptide levels during wound healing are difficult to compare due to vastly differing experimental models and neuropeptides of interest. Jonsson *et al* showed an immediate increase in SP immunoreactivity in

dog paw lymph after scalding injury, indicating release of SP during wounding [178]. Grönblad *et al* observed many more SP and CGRP nerve fibers in healing rabbit medial collateral ligaments as compared to normal ligaments, suggesting involvement of these nerves in healing [179]. Senapati *et al*, however, observed significant depletion of SP, CGRP, and somatostatin in rat skin [180] within 1 d of wounding that persisted for 14 d, consistent with release of neuropeptides at the wound site but also possibly explained by decreased synthesis and transport to the nerve terminals. These data, which support the ideas of an increase in neuropeptidergic tone at the wound site, are further substantiated by studies in which neuropeptide replacement enhanced wound repair. In a preliminary study, Spevak *et al* found that wounds in rat skin healed faster when treated with SP [181]. English *et al* demonstrated a trophic action of the CSNS on wound healing by showing that inclusion of touch domes in skin grafts enhanced the return of neural function in the graft and, most importantly, the quality and survivability of the graft [182,183]. (Touch domes are specialized areas of skin that are richly innervated and are known to provide tropic or trophic factors for regenerating nerves [184]).

CONCLUSIONS

There is increasing evidence that the nervous system can modulate tissue inflammatory, proliferative, and reparative processes. The role of the CSNS and neuropeptides such as SP in inflammation and tissue repair is determined not only by the synthesis and secretion of neuropeptides such as SP, but also by availability of and interaction with its receptor on target cells and by the rate of SP degradation by tissue peptidases such as NEP. Furthermore, the rate at which reinnervation of wounds occurs may have dramatic effects on the time course and quality of tissue repair, thus pointing toward a crucial role for neurotropic and neurotrophic factors like NGF.

Our current understanding of the different components of this complex system suggests that defects in skin integrity and wound healing in patients with neurologic sensory deficits may be caused by abnormalities in growth and function of the CSNS. In view of the scale of the clinical problem represented by chronic wounds, increasing our knowledge of the interaction between the CSNS and wound healing has the potential for making a significant impact on therapy.

REFERENCES

- Bernstein J: Neuropeptides and the skin. In: Goldsmith L (ed.). *Physiology, Biochemistry, and Molecular Biology of the Skin*. Oxford University Press, New York, 1991, pp 816-835
- Eedy D: Neuropeptides in skin. *Br J Dermatol* 128:597-605, 1993
- Hosoi J, Murphy G, Egan C, Lerner E, Grabbe S, Asahina A, Granstein R: Regulation of Langerhans cell function by nerves containing calcitonin gene-related peptide. *Nature* 363:159-163, 1993
- Baraniuk J, Marek L, Kowalski L, Kaliner M: Neuropeptides in the skin. In: Bos J (ed.). *Skin Immunology*. CRC Press, Boca Raton, 1990, pp 307-326
- Pincelli C, Rantini F, Giannetti A: Neuropeptides and skin inflammation. *Dermatology* 187:153-158, 1993
- Holzer P: Local effector functions of capsaicin-sensitive sensory nerve endings: involvement of tachykinins, calcitonin gene-related peptide and other neuropeptides. *Neuroscience* 24:739-768, 1988
- Chapman L, Goodell H: The participation of the nervous system in the inflammatory reaction. *Ann NY Acad Sci* 116:990-1017, 1964
- Jancsó N, Jancsó-Gábor A, Szolcsányi J: Direct evidence for neurogenic inflammation and its prevention by denervation and by pretreatment with capsaicin. *Br J Pharmacol* 31:138-51, 1967
- Foreman J, Jordan C: Neurogenic inflammation. *Trends Pharmacol Sci* 5:116-119, 1984
- Nilsson J, von Euler A, Dalsgaard C-J: Stimulation of connective tissue cell growth by substance P and substance K. *Nature* 315:61-63, 1985
- Tanaka T, Danno K, Ikai K, Imamura S: Effects of substance P and substance K on the growth of cultured keratinocytes. *J Invest Dermatol* 90:399-401, 1988
- Skofitsch G, Jacobowitz D: Calcitonin gene-related peptide coexists with substance P in capsaicin sensitive neurons and sensory ganglia of the rat. *Peptides* 6:747-754, 1985
- Brain S, Williams T, Tippins J, Morris H, MacIntyre I: Calcitonin gene-related peptide is a potent vasodilator. *Nature* 313:54-56, 1985
- Haegerstrand A, Dalsgaard C-J, Jonzon B, Larsson O, Nilsson J: Calcitonin gene-related peptide stimulates proliferation of human endothelial cells. *Proc Natl Acad Sci USA* 87:3299-3303, 1990

15. Dalsgaard C-J, Hägerstrand A, Theodorsson-Norheim E, Brodin E, Hökfelt T: Neurokinin A-like immunoreactivity in rat primary sensory neurons: coexistence with substance P. *Histochemistry* 83:37-39, 1985
16. Nilsson J, Sejersen T, Nilsson A, Dalsgaard C-J: DNA synthesis induced by the neuropeptide substance K correlates to the level of myc-gene transcripts. *Biochem Biophys Res Commun* 137:167-174, 1986
17. Williams T: Vasoactive intestinal polypeptide is more potent than prostaglandin E₂ as a vasodilator and oedema potentiator in rabbit skin. *Br J Pharmacol* 77:505-509, 1982
18. Leah H, Cameron A, Kelly W, Snow P: Coexistence of peptide immunoreactivity in sensory neurons of the cat. *Neuroscience* 16:683-690, 1985
19. Haegerstrand A, Jonzon B, Dalsgaard C-J, Nilsson J: Vasoactive intestinal polypeptide stimulates cell proliferation and adenylyl cyclase activity of cultured human keratinocytes. *Proc Natl Acad Sci USA* 86:5993-5996, 1989
20. Rabier M, Farber E, Wilkinson D: Neuropeptides modulate leukotriene B₄ mitogenicity toward cultured human keratinocytes. *J Invest Dermatol* 100:132-136, 1993
21. Barber R, Vaughn J, Slemmon J, Salvaterra P, Roberts E, Leeman S: The origin, distribution and synaptic relationships of substance P axons in rat spinal cord. *J Comparative Neurol* 184:331-351, 1979
22. Ulrich A, Gray A, Berman C, Dull T: Human beta-nerve growth factor gene sequence is highly homologous to that of mouse. *Nature* 303:821-825, 1983
23. Diamond J, Holmes M, Coughlin M: Endogenous NGF and nerve impulses regulate the collateral sprouting of sensory axons in the skin of the adult rat. *J Neurosci* 12:1454-1466, 1992
24. Kalchauer C, Barde Y-A, Thoenen H, Le Douarin N: *In vivo* effect of brain-derived neurotrophic factor on the survival of developing dorsal root ganglion cells. *EMBO J* 6:2871-2873, 1987
25. Leibrock J, Lottspeich F, Hohn A, et al: Molecular cloning and expression of brain-derived neurotrophic factor. *Nature* 341:149-152, 1989
26. Ernors P, Ibáñez C, Ebendal T, Olson L, Persson H: Molecular cloning and neurotrophic activities of a protein with structural similarities to nerve growth factor: developmental and topographical expression in the brain. *Proc Natl Acad Sci USA* 87:5454-5458, 1990
27. Berkemeier L, Winslow J, Kaplan D, Nikolics K, Goeddel D, Rosenthal A: Neurotrophin-5: a novel neurotrophic factor that activates trk and trkB. *Neuron* 7:857-866, 1991
28. Tron V, Coughlin M, Jang D, Stanis J, Sauder D: Expression and modulation of nerve growth factor in murine keratinocytes (PAM 212). *J Clin Invest* 85:1085-1089, 1990
29. Di Marco E, Marshisio P, Bondanza S, Franzi A, Cancedda R, De Luca M: Growth-regulated synthesis and secretion of biologically active nerve growth factor by human keratinocytes. *J Biol Chem* 266:21718-21722, 1991
30. Hökfelt T, Johansson O, Ljungdahl A, Lundberg J, Schultzberg M: Peptidergic neurons. *Nature* 284:515-521, 1980
31. Pernow B: Substance P. *Pharmacol Rev* 35:85-141, 1983
32. Pickel V, Reis D, Leeman SE: Ultrastructural localization of substance P in neurons of rat spinal cord. *Brain Res* 122:534-540, 1977
33. Nawa H, Kotani H, Nakanishi S: Tissue-specific generation of two prepro-tachykinin mRNAs from one gene by alternative RNA splicing. *Nature* 312:729-734, 1984
34. Lynch D, Snyder S: Neuropeptides: multiple molecular forms, metabolic pathways, and receptors. *Annu Rev Biochem* 55:773-799, 1986
35. Tatemoto K, Lundberg J, Jörmvall H, Mutt V: Neuropeptide K: isolation, structure and biological activities of a novel brain tachykinin. *Biochem Biophys Res Commun* 128:947-953, 1985
36. Nakanishi S: Substance P precursor and Kininogen: their structures, gene organizations, and regulation. *Physiol Rev* 67:1117-1142, 1987
37. Burcher E, Buck SH: Multiple binding sites in hamster, rat, and guinea-pig urinary bladder. *Eur J Pharmacol* 1986:128:165-177
38. Muramatsu I, Nakanishi S, Fujiwara M: Comparison of the responses to the sensory neuropeptides, substance P, neurokinin A, neurokinin B and calcitonin gene-related peptide and to trigeminal nerve stimulation in the iris sphincter muscle of the rabbit. *Jpn J Pharmacol* 44:85-92, 1987
39. Roques B, Fournie-Zaluski M, Soroica E, Lecomte J, Malfroy B, Llorens C: The enkephalinase inhibitor thiorphan shows antinociceptive activity in mice. *Nature* 288:286-288, 1980
40. Matsas R, Fulcher I, Kenny A, Turner A: Substance P and [Leu]enkephalin are hydrolyzed by an enzyme in pig caudate synaptic membranes that is identical with the endopeptidase of kidney microvilli. *Proc Natl Acad Sci USA* 80:3111-3115, 1983
41. Matsas R, Kenny A, Turner A: The metabolism of neuropeptides. The hydrolysis of peptides, including enkephalins, tachykinins and their analogues. *Biochem J* 1984:223:433-440, 1984
42. Hershey A, Krause J: Molecular characterization of a functional cDNA encoding the rat substance P receptor. *Science* 247:958-962, 1990
43. Hershey A, Dykema P, Krause J: Organization, structure, and expression of a gene encoding the rat substance P receptor. *J Biol Chem* 266(7):4366-4374, 1991
44. Takeda Y, Chou K, Taked J, Sachais B, Krause J: Molecular cloning, structural characterization and functional expression of the human substance P receptor. *Biochem Biophys Res Commun* 179:1232-1240, 1991
45. Sasaki Y, Nakanishi S: Molecular characterization of rat substance K receptor and its mRNAs. *Biochem Biophys Res Commun* 165:695-702, 1989
46. Gerard N, Edy RJ, Shows T, Gerard C: The human neurokinin A (substance K) receptor. Molecular cloning of the gene, chromosomal localization, and isolation of cDNA from tracheal and gastric tissues. *J Biol Chem* 265:20455-20462, 1990
47. Maggio J: Tachykinins. *Annu Rev Neurosci* 11:13-28, 1988
48. Quirion R, Dam T: Multiple neurokinin receptors: recent developments. *Regul Pept* 22:18-25, 1988
49. Helke C, Krause J, Mantyh P, Couture R, Bannon M: Diversity in tachykinin peptidergic neurons: multiple peptides, receptors and regulatory mechanisms. *FASEB J* 4:1606-1615, 1990
50. Tsuchida K, Shigemoto R, Yokota Y, Nakanishi S: Tissue distribution and quantitation of the mRNAs for three rat tachykinin receptors. *Eur J Biochem* 193:751-757, 1990
51. Buck S, Helke C, Burcher E, Shults C, O'Donohue T: Pharmacologic characterization and autoradiographic distribution of binding sites for iodinated tachykinins in the rat central nervous system. *Peptides* 7:1109-1120, 1986
52. Saffroy M, Beaujouan J-C, Torrens Y, Besseyre J, Bergström L, Glowinski J: Localization of tachykinin binding sites (NK1, NK2, NK3 ligands) in the rat brain. *Peptides* 9:227-241, 1988
53. Mantyh P, Gates T, Mantyh C, Maggio J: *J Neurosci* 9:258-279, 1989
54. Hershey A, Plenzani L, Woodward R, Miledi R, Krause J: Molecular and genetic characterization, functional expression, and mRNA expression patterns of a rat substance P receptor. *Ann NY Acad Sci* 632:63-78, 1991
55. Krause J, Takeda Y, Hershey A: Structure, functions, and mechanisms of substance P receptor action. *J Invest Dermatol* 98:2S-7S, 1992
56. Ansel J, Brown J, Payan D, Brown M: Substance P selectively activates TNF- α gene expression in murine mast cells. *J Immunol* 150:1-8, 1993
57. Brown J, Perry P, Hefeneider S, Ansel J: Neuropeptide modulation of keratinocyte cytokine production. In: Oppenheim, Powanda, Kluger, Dinarello (eds.), *Molecular and Cellular Biology of Cytokines*. Wiley-Liss, Inc., 1990, pp 451-456
58. Matis W, Lavker R, Murphy G: Substance P induces the expression of an endothelial-leukocyte adhesion molecule by microvascular endothelium. *J Invest Dermatol* 94:492-495, 1990
59. Greeno E, Mantyh P, Vercellotti G, Moldow C: Functional neurokinin 1 receptors for substance P are expressed by human vascular endothelium. *J Exp Med* 45:1269-1276, 1992
60. Phillis J, Limacher J: Substance P excitation of cerebral cortical Betz cells. *Brain Res* 69:158-163, 1974
61. Rivier C, Brown M, Vale W: Effect of neurotensin, substance P and morphine sulfate on the secretion of prolactin and growth hormone in the rat. *Endocrinology* 100:751-754, 1977
62. Vijayan E, McCann S: *In vivo* and *in vitro* effects of substance P and neurotensin on gonadotropin and prolactin release. *Endocrinology* 105:64-68, 1979
63. Eckstein N, Wehrenberg W, Louis K, Carmel P, Zimmermann E, Frantz A: Effects of substance P on anterior pituitary secretion in the female rhesus monkey. *Neuroendocrinology* 31:338-342, 1980
64. Arisawa M, De Palatis L, Ho R, et al: Stimulatory role of substance P on gonadotropin release in ovariectomized rats. *Neuroendocrinology* 51:523-529, 1990
65. Traczysk W, Pau K-YF, Kaynard A, Spies H: Modulatory role of substance P on gonadotropin and prolactin secretion in the rabbit. *J Physiol Pharmacol* 43:279-288, 1992
66. Hedner T, Hedner J, Jonason J, Lundberg D: Evidence suggesting a role for substance P in central respiratory regulation in the rat. *Acta Physiologica Scandinavica* 1981:112:487-489, 1981
67. Yamamoto Y, Lagercrantz H, von Euler C: Effects of substance P and TRH on ventilation and pattern of breathing in newborn rabbits. *Acta Physiologica Scandinavica* 113:541-543, 1981
68. Rossell S, Björkroth U, Chang D, et al. In: von Euler U, Pernow B (eds.), *Substance P*. Raven Press, New York, 1977, p 83
69. Bury R, Mahford M: Interactions between local anesthetics and spasmogens on the guinea-pig ileum. *J Pharmacol Exp Ther* 197:633-640, 1976
70. Lundberg J, Saria A, Brodin E, Rosell S, Folkers K: A substance P antagonist inhibits vagally induced increase in vascular permeability and bronchial smooth muscle contraction in the guinea pig. *Proc Natl Acad Sci USA* 80:1120-1124, 1983
71. Lundberg J, Saria A: Bronchial smooth muscle contraction induced by stimulation of capsaicin-sensitive sensory neurons. *Acta Physiologica Scandinavica* 116:473-476, 1982
72. Lundberg J, Saria A: Capsaicin-induced desensitization of airway mucosa to cigarette smoke, mechanical and chemical irritants. *Nature* 302:251-253, 1983
73. Bernstein J, Hamill J: Substance P. *J Invest Dermatol* 77:250-251, 1981
74. Borson D, Corrales R, Varsano S, Gold M, Viro N, Caughey G: Enkephalinase inhibitors potentiate substance P-induced secretion of ³⁵SO₄-macromolecules from ferret trachea. *Exp Lung Res* 12:21-36, 1987
75. Liang T, Cascieri M: Substance P stimulation of amylase release by isolated parotid cells and inhibition of substance P induction of salivation by vasoactive peptides. *Mol Cell Endocrinol* 15:151-162, 1979
76. Sjödin L, Conlon T, Gustavson C, Uddholm K: Interaction of substance P with dispersed pancreatic acinar cells from the guinea pig. *Acta Physiologica Scandinavica* 120:21-26, 1984
77. Brown M, Vale W: Effects of neurotensin and substance P on plasma insulin, glucagon, and glucose levels. *Endocrinology* 98:819-822, 1976
78. Kaneto A, Kaneko T, Kajinuma H, Kosaka K: Effects of substance P and neurotensin infused intrapancreatically on glucagon and insulin secretion. *Endocrinology* 102:393-401, 1978
79. Lundquist I, Sundler F, Ahren B, Alumets J, Hakanson R: Somatostatin, pancreatic polypeptide, substance P, and neurotensin: cellular distribution and

- effects on stimulated insulin secretion in the mouse. *Endocrinology* 104:832-838, 1979
80. Skrabanek P, Cannon D, Kirrane J, Legge D, Powell D: Circulating immunoreactive substance P in man. *Irish J Med Sci* 145:399-408, 1976
 81. Powell D, Skrabanek P, Cannon D: In: von Euler U, Pernow B (eds.). *Substance P*. Raven Press, New York, 1977, p 35
 82. Powell D, Skrabanek P, Cannon D, Balfé A: In: Marsan C, Traczyk W (eds.). *Neuropeptides and Neural Transmissions*. Raven Press, New York, 1980, p 105
 83. Pernow B, Rosell S: Effect of substance P on blood flow in canine adipose tissue and skeletal muscle. *Acta Physiologica Scandinavica* 93:139-141, 1975
 84. Burcher E, Atterhog J, Pernow B, Rosell S: In: von Euler U, Pernow B (eds.). *Substance P*. Raven Press, New York, 1977, p 261
 85. Eklund B, Jogenstrand T, Pernow B: In: von Euler U, Pernow B (eds.). *Substance P*. Raven Press, New York, 1977, p 275
 86. Hokfelt T, Kellerth J, Nilsson G, Pernow B: Substance P: localization in the central nervous system and in some primary sensory neurons. *Science* 190:889-890, 1975
 87. Bolton T, Clapp L: Endothelial-dependent relaxant actions of carbachol and substance P in arterial smooth muscle. *Br J Pharmacol* 87:713-715, 1986
 88. Cuella A, Del Fiaco M, Paxinos G: The central and peripheral ends of the substance P-containing sensory neurons in the rat trigeminal system. *Brain Res* 152:499-500, 1978
 89. Bill A, Stjernschantz J, Mandahl A, Brodin E, Nilsson G: Substance P: release on trigeminal nerve stimulation, effects in the eye. *Acta Physiologica Scandinavica* 106:371-373, 1979
 90. Olgart L, Gazelius B, Brodin E, Nilsson G: Release of substance P-like immunoreactivity from the dental pulp. *Acta Physiologica Scandinavica* 101:510-512, 1977
 91. Lembeck F, Holzer P: Substance P as neurogenic mediator of antidromic vasodilation and neurogenic plasma extravasation. *Naunyn-Schmiedeberg's Arch Pharmacol* 310:175-183, 1979
 92. Erjavec F, Lembeck F, Florjanc-Imman T, et al: Release of histamine by substance P. *Naunyn-Schmiedeberg's Arch Pharmacol* 317:67-70, 1981
 93. Coutts A, Jorizzo J, Greaves M, Burnstock G: Mechanism of vasodilatation due to substance P in humans. *Br J Dermatol* 105:354-355, 1981
 94. Devillier P, Regoli D, Asseraf A, Descours B, Marsac J, Renoux M: Histamine release and local responses of rat and human skin to substance P and other mammalian tachykinins. *Pharmacology* 32:340-347, 1986
 95. Hagermark O, Hokfelt T, Pernow B: Flare and itch induced by substance P in human skin. *J Invest Dermatol* 71:233-235, 1978
 96. Barnes P, Brown M, Dollery C, Fuller R, Heavey D, Ind P: Histamine is released from skin by substance P but does not act as the final vasodilator in the axon reflex. *Br J Pharmacol* 88:741-745, 1986
 97. Foreman J, Jordan C, Oehme P, Renner H: Structure-activity relationships for some substance P-related peptides that cause wheal and flare reactions in human skin. *J Physiol* 335:449-465, 1983
 98. Payan D, Brewster D, Goetzl E: Specific stimulation of human T lymphocytes by substance P. *J Immunol* 131:1613-1615, 1983
 99. Stanis A, Befus D, Bienenstock J: Differential effect of vasoactive intestinal polypeptide, substance P, and somatostatin on immunoglobulin synthesis and proliferation by lymphocytes from Peyer's patches, mesenteric lymph nodes, and spleen. *J Immunol* 136:152-156, 1987
 100. Scicchitano R, Bienenstock J, Stanis A: In vivo immunomodulation by the neuropeptide substance P. *Immunology* 63:733-735, 1988
 101. Levine J, Clark R, Devor M, Helms C, Moskowitz M, Basbaum A: Intraneuronal substance P contributes to the severity of experimental arthritis. *Science* 226:547-549, 1984
 102. Pascual D, Xu-Amano J, Kiyono H, McGee J, Bout K: Substance P acts directly upon cloned B lymphoma cells to enhance IgA and IgM production. *J Immunol* 146:2130-2136, 1991
 103. McGillis J, Mitsuhashi M, Payan D: Immunologic properties of substance P. In: Ader R, Felten D, Cohen N (eds.). *Psychoneuroimmunology*. Academic Press, San Diego, 1991, pp 209-223
 104. Sudduth-Klinger J, Gilbert M, Bunnett N, Payan D: Characterization of a stable lymphocyte cell line transfected with the substance P receptor cDNA (submitted)
 105. Calvo C-F, Chavanel G, Senik A: Substance P enhances IL-2 expression in activated human T cells. *J Immunol* 148:3498-3504, 1992
 106. Payan D, Levine J, Goetzl E: Modulation of immunity and hypersensitivity by sensory neuropeptides. *J Immunol* 132:1601-1604, 1984
 107. Baxter J, Adamik R: Differences in requirements and actions of various histamine-releasing agents. *Biochem Pharmacol* 27:497-503, 1978
 108. Hartung H, Toyka K: Activation of macrophages by substance P: induction of oxidative burst and thromboxane release. *Eur J Pharmacol* 89:301-305, 1983
 109. Payan D: Receptor-mediated mitogenic effects of substance P on cultured smooth muscle cells. *Biochem Biophys Res Commun* 130:104-109, 1985
 110. Ziche M, Morbidelli L, Pacini M, Geppetti P, Alesandri G: Substance P stimulated neovascularization in vivo and proliferation of cultured endothelial cells. *Microvascular Res* 40:264-266, 1990
 111. Silver A: *The Biology of Cholinesterases*. Elsevier, Amsterdam, 1974
 112. Iversen L: In: Iversen L, Iversen S, Synder S (eds.). *Handbook of Psychopharmacology*, Vol. 3. Plenum Press, New York, 1978, pp 381-442
 113. Bunnett N: Release and breakdown. Postsecretory metabolism of peptides. *Am Rev Respiratory Dis* 136:S27-S34, 1987
 114. Wong-Leung Y, Kenny A: Some properties of a microsomal peptidase in rat kidney. *Biochem J* 110:5P, 1968
 115. Kerr M, Kenny A: The purification and specificity of a neutral endopeptidase from rabbit kidney brush border. *Biochem J* 137:477-488, 1974
 116. Kerr M, Kenny A: The molecular weight and properties of a neutral metallo-endopeptidase from rabbit kidney brush border. *Biochemical J* 137:489-495, 1974
 117. Malfroy B, Swerts J, Guyon A, Roques B, Schwartz J: High-affinity enkephalin-degrading peptidase in brain is increased after morphine. *Nature* 276:523-526, 1978
 118. Matsas R, Kenny A, Turner A: The metabolism of neuropeptides. The hydrolysis of peptides, including enkephalins, tachykinins, and their analogs, by endopeptidase 24.11. *Biochem J* 223:433-440, 1984
 119. Turner A: Chapter 10. In: Turner A (ed.). *Neuropeptides and Their Peptidases*. VCH Publishers, New York, 1987, pp 183-201
 120. Bunnett N, Turner A, Hryszko J, Kobayashi R, Walsh J: Isolation of endopeptidase-24.11 (EC3.4.24.11, "enkephalinase") from the pig stomach. *Gastroenterology* 95:952-957, 1988
 121. de la Baume S, Gros C, Yi C, et al: Selective participation of both "enkephalinase" and aminopeptidase activities in the metabolism of endogenous enkephalins. *Life Sci* 31:1753-1766, 1982
 122. Waksman G, Bouboutou R, Devin J, et al: In vitro and in vivo effects of ketorphan on enkephalin metabolism in rodent brain. *Eur J Pharmacol* 117:233-243, 1985
 123. Cailliet P, Marçais-Collado H, Costentin J, Yi C, de la Baume S, Schwartz J: Inhibition of enkephalin metabolism by, and antinociceptive activity of, bestatin, an aminopeptidase inhibitor. *Eur J Pharmacol* 86:329-336, 1983
 124. Erdos E, Skidgel R: Neutral endopeptidase 24.11 (enkephalinase) and related regulators of peptide hormones. *FASEB J* 3:145-151, 1989
 125. Duser D, Umeno E, Graf P, Djokic T, Borson D, Nadel J: Airway neutral endopeptidase-like enzyme modulates tachykinin-induced bronchoconstriction in vivo. *J Appl Physiol* 65:2585-2591, 1988
 126. Kohrogi H, Graf P, Sekizawa K, Borson D, Nadel J: Neutral endopeptidase inhibitors potentiate substance P- and capsaicin-induced cough in awake guinea pigs. *J Clin Invest* 82:2063-2068, 1988
 127. Umeno E, Nadel J, Huang H-T, McDonald D: Inhibition of neutral endopeptidase potentiates neurogenic inflammation in the rat trachea. *J Appl Physiol* 66:2647-2652, 1989
 128. Shipp M, Tarr G, Chen C-Y, et al: CD10/neutral endopeptidase 24.11 hydrolyzes bombesin-like peptides and regulates the growth of small cell carcinomas of the lung. *Proc Natl Acad Sci USA* 88:10662-10666, 1991
 129. Shipp M, Stefano G, D'Adamo L, et al: Down-regulation of enkephalin-mediated inflammatory responses by CD10/neutral endopeptidase 24.11. *Nature* 347:394-396, 1991
 130. Borson D, Brokaw J, Sekizawa K, McDonald D, Nadel J: Neutral endopeptidase and neurogenic inflammation in rats with respiratory infections. *J Appl Physiol* 66:2653-2658, 1989
 131. Hwang L, Leichter R, Okamoto A, Payan D, Collins S, Bunnett N: Down-regulation of neutral endopeptidase (EC 3.4.24.11) in the inflamed rat intestine. *Am J Physiol* 264:G735-G743, 1993
 132. Borson D, Gruenert D: Glucocorticoids induce neutral endopeptidase in transformed human tracheal epithelial cells. *Am J Physiol* 260:L83-L89, 1991
 133. Ihara H, Nakanishi S: *J Biol Chem* 265:22441-22445, 1990
 134. Cohen S, Levi-Montalcini R, Hamburger V: A nerve growth stimulating factor isolated from sarcoma 37 and 180. *Proc Natl Acad Sci USA* 40:1014-1018, 1954
 135. Levi-Montalcini R: The nerve growth factor: thirty-five years later. *EMBO J* 6:1145-1154, 1987
 136. Greene L, Shooter E: The nerve growth factor: biochemistry, synthesis, and mechanism of action. *Annu Rev Neurosci* 3:353-402, 1980
 137. Heumann R, Korsching S, Scott J, Thoenen H: Relationship between level of nerve growth factor (NGF) and its messenger RNA in sympathetic ganglia and peripheral target tissues. *EMBO J* 3:3183-3189, 1984
 138. Whittemore S, Ebendal T, Larkfors L, et al: Developmental regional expression of β nerve growth factor messenger RNA and protein in the rat central nervous system. *Proc Natl Acad Sci USA* 83:817-821, 1986
 139. Yaar M, Grossman K, Eller M, Gilchrist B: Evidence for nerve growth factor-mediated paracrine effects in human epidermis. *J Cell Biol* 115:821-828, 1991
 140. Sutter A, Riopelle R, Harris-Warrick R, Shooter E: Nerve growth factor receptors: characterization of two distinct classes of binding sites on chick embryo sensory ganglion cells. *J Biol Chem* 254:5972-5982, 1979
 141. Bothwell M: Keeping track of neurotrophin receptors. *Cell* 65:915-918, 1991
 142. Klein R, Jing S, Nandori V, O'Rourke E, Barbacid M: The *trk* proto-oncogene encodes a receptor to nerve growth factor. *Cell* 65:189-197, 1991
 143. Kaplan D, Hempstead B, Martin-Zanca D, Chao M, Parada L: The *trk* protooncogene product: a signal transducing receptor for nerve growth factor. *Science* 252:554-558, 1991
 144. Levi-Montalcini R, Angeletti P: Essential role of the nerve growth factor in the survival and maintenance of dissociated sensory and sympathetic embryonic nerve cells in vitro. *Dev Biol* 7:653-659, 1963
 145. Johnson E, Gorin P, Bandeis L, Pearson J: Dorsal root ganglion neurons are destroyed by exposure in utero to maternal antibodies to nerve growth factor. *Science* 210:916-918, 1980
 146. Ross M, Lofstrand S, Gorin P, Johnson E, Schwartz J: Use of an experimental autoimmune model to define nerve growth factor dependency of peripheral and central substance P containing neurons in the rat. *J Neurosci* 1:1304-1311, 1981
 147. Stockel K, Schwab M, Thoenen H: Specificity of retrograde transport of nerve growth factor (NGF) in sensory neurons: a biochemical and morphological study. *Brain Res* 89:1-14, 1975
 148. Schwartz J, Pearson J, Johnson E: Effect of exposure to anti-NGF on sensory neurons of adult rats and guinea pigs. *Brain Res* 244:378-381, 1982

149. Waris T: Reinnervation of free skin autografts in the rat. *Plastic Reconstruct Surg* 12:85-93, 1978
150. Waris T: Degeneration and regeneration of nerves in a dorsal skin flap in the rat. *Scand J Plastic Reconstruct Surg Hand Surg* 12:95-103, 1978
151. Kishimoto S, Maruo M, Ohse C, et al: The regeneration of the sympathetic catecholaminergic nerve fibers in the process of burn wound healing in guinea pigs. *J Invest Dermatol* 79:141-146, 1982
152. Kishimoto S: The regeneration of substance P-containing nerve fibers in the process of burn wound healing in the guinea pig skin. *J Invest Dermatol* 83:219-223, 1984
153. Aldskogius H, Hermanson A, Jonsson C-E: Reinnervation of experimental superficial wounds in rats. *Plastic Reconstruct Surg* 79:595-599, 1987
154. Devor M, Schonfeld D, Seltzer Z, Wall P: Two modes of cutaneous reinnervation following peripheral nerve injury. *J Compar Neurol* 185:211-220, 1979
155. Owen D, Logan A, Robinson P: A role for nerve growth factor in collateral reinnervation from sensory nerves in the guinea pig. *Brain Res* 476:156-160, 1989
156. Doubleday B, Robinson P: The role of nerve growth factor in collateral reinnervation by cutaneous C-fibers in the rat. *Brain Res* 593:179-184, 1992
157. Wilson P, Scott J, Penschow J, Coghlan J, Rush R: Identification and quantification of mRNA for nerve growth factor in histological preparations. *Neurosci Lett* 64:323-329, 1986
158. Taniuchi M, Clark H, Johnson E: Induction of nerve growth factor receptor in Schwann cells after axotomy. *Proc Natl Acad Sci USA* 83:4094-4098, 1986
159. Assouline J, Bosch P, Lim R, Kim I, Jensen R, Pantazis N: Rat astrocytes and Schwann cells in culture synthesize nerve growth factor like neurite-promoting factors. *Dev Brain Res* 31:103-118, 1987
160. Taniuchi M, Clark H, Schucitzer J, Johnson EJ: Expression of nerve growth factor receptors by Schwann cells of axotomized peripheral nerves: ultrastructural location, suppression by axonal contact binding properties. *J Neurosci* 8:664-681, 1988
161. Diamond J, Foerster A, Holmes M, Coughlin M: Sensory nerves in adult rats regenerate and restore sensory function to the skin independently of endogenous NGF. *J Neurosci* 12:1467-1476, 1992
162. Pearce F, Thompson H: Some characteristics of histamine secretion from rat peritoneal mast cells stimulated with nerve growth factor. *J Physiol* 372:379-393, 1986
163. Matsuda H, Coughlin M, Bienenstock J, Denburg J: Nerve growth factor promotes human hemopoietic colony growth and differentiation. *Proc Natl Acad Sci USA* 85:6508-6512, 1988
164. Otten U, Ehrhard P, Peck R: Nerve growth factor induces growth and differentiation of human B lymphocytes. *Proc Natl Acad Sci USA* 86:10059-10063, 1989
165. Lewis T, Marvin H: Observations relating to vasodilatation arising from antidromic impulses, to herpes zoster and trophic effects. *Heart* 14:27-46, 1927
166. Bayliss W: On the origin from the spinal cord of vasodilator fibers of the hind limb, and on the nature of these fibers. *J Physiol* 26:173-209, 1901
167. Lewis T: The nocifensor system of nerves and its reactions. *Br Med J* 1:431-435, 1927
168. Johnson P, Doll S: Dermal nerves in human diabetic subjects. *Diabetes* 33:244-250, 1984
169. Jancsó G, Obál F, Tóth-Kasa I, Katona M, Husz S: The modulation of cutaneous inflammatory reactions by peptide-containing sensory nerves. *Int J Tissue Reactions* 7:449-457, 1985
170. Olivari N, Schrudde J, Wahle H: The surgical treatment of bedsores in paraplegics. *Plastic Reconstruct Surg* 50:477-482, 1972
171. Basson M, Burney R: Defective wound healing in patients with paraplegia and quadriplegia. *Surg Gynecol Obstet* 155:9-12, 1982
172. Lusthaus S, Shoshan S, Benmeir P, Livoff A, Ashur H, Vardy D: Effect of denervation on incision wound scars in rabbits. *J Geriatr Dermatol* 1:11-14, 1993
173. Nagy J, Iversen L, Goedert M, Chapman D, Hunt S: Dose-dependent effects of capsaicin on primary sensory neurons in the neonatal rat. *J Neurosci* 3:399-406, 1983
174. Kjartansson J, Dalsgaard C-J, Jonsson C-E: Decreased survival of experimental critical flaps in rats after sensory denervation with capsaicin. *Plastic Reconstruct Surg* 79:218-221, 1987
175. Shimizu T, Fujita S, Izumi K, Koja T, Ohba N, Fukuda T: Corneal lesions induced by the systemic administration of capsaicin in neonatal mice and rats. *Naunyn-Schmiedeberg's Arch Pharmacol* 326:347-351, 1984
176. Duke-Elder S, Leigh A: Diseases of the outer eyes. In: Duke-Elder S (ed.). *System of Ophthalmology*. Henry Kimble, London, 1965, pp 803-810
177. Gallar J, Pozo M, Rebello I, Belmonte C: Effects of capsaicin on corneal wound healing. *Invest Ophthalmol Vis Sci* 31:1968-1974, 1990
178. Jonsson C-E, Brodin E, Dalsgaard C-J, Haegerstrand A: Release of substance-P-like immunoreactivity in dog paw lymph after scalding injury. *Acta Physiol Scand* 126:21-24, 1986
179. Grönlund M, Korkala O, Kontinen Y, Kuokkanen H, Liesi P: Immunoreactive neuropeptides in nerves in ligamentous tissue. An experimental neuroimmunohistochemical study. *Clin Orthopaed Relat Res* 265:291-296, 1991
180. Senapati A, Anand P, McGregor G, Ghatei M, Thompson R, Bloom S: Depletion of neuropeptides during wound healing in rat skin. *Neurosci Lett* 71:101-105, 1986
181. Spevak S, Shekhter A, Hilse H, Oehme P, Soloveva A: Aspects of wound healing in spontaneous hypertensive rats. The effect of morphine, substance P and its fragment SP₁₋₄. *Biull Eksp Biol Med* 1989;107(6):739-43
182. English K, Stayner N, Krueger G, Tuckett R: Tactile function in skin-equivalent grafts. *Exp Neurol* 115:104-108, 1992
183. English K, Stayner N, Krueger G, Tuckett R: Functional innervation of cultured skin grafts. *J Invest Dermatol* 99:120-128, 1992
184. Horch K: The influence of mechanoreceptor structures on regenerating sensory axons after cutaneous nerve transection in the cat. *Neurosci Lett* 32:281-284, 1982
185. Hartung H, Wolters K, Toyka K: Substance P: binding properties and studies on cellular responses in guinea pig macrophages. *J Immunol* 136:3856-3863, 1986
186. Wiederman C, Wiederman F, Apperl A, Kieselbach G, Konwalinka G, Braunsteiner H: In vitro human polymorphonuclear leukocyte chemokinesis and human monocyte chemotaxis are different activities of aminoterminal and carboxyterminal substance P. *Arch Pharmacol* 340:185-190, 1989
187. Ruff M, Whal S, Pert C: Substance P receptor-mediated chemotaxis of human monocytes. *Peptides* 6(suppl):107-111, 1985
188. Sudduth-Klinger J, Schumann M, Gardner P, Payan D: Functional and immunological responses of Jurkat lymphocytes transfected with the substance P receptor. *Cell Mol Neurobiol* 12:379-395, 1992